

Cannock Chase District Nature Recovery Network Mapping



FINAL REPORT

Staffordshire Wildlife Trust 2020



Table of Contents

1. Executive Summary	3
2. Introduction	4
3. Review of previous biodiversity opportunity mapping assessments	6
4. Existing evidence base review	7
4.1 Available environmental datasets.....	7
4.2 Phase 1 habitat mapping via aerial photography interpretation	7
4.3 National Character Areas in Cannock Chase District.....	8
4.4 Minerals Safeguarding Zones in Cannock Chase District.....	9
4.5 Data used and limitations	10
5. Mapping the opportunities to enhance habitats for biodiversity	11
5.1 Habitat distinctiveness mapping.....	11
5.2 Habitat distinctiveness mapping limitations	14
5.3 Biodiversity metric 2.0	14
5.4 Strategic Habitat Areas	15
6. Establishing the Habitat Connectivity Opportunity Areas (HCO) for Cannock Chase District	18
6.1 Habitat Connectivity Opportunity Areas Rationale	18
7. Results	19
7.1 Habitat Connectivity Opportunity Areas identified	19
7.2 Woodland Opportunity Area	21
7.3 Grassland Opportunity Area	25
7.4 Heathland Opportunity Area	28
7.5 Wetland Opportunity Area	31
7.6 Pasture and Arable Opportunity Area	35
7.7 Urban Fabric Opportunity Area	38
8. Cross boundary habitat connectivity	42
9. Practical Application of the maps.....	44
10. Next Steps	45
10.1 Habitat connectivity bottlenecks	45
10.2 How the strategic mapping will evolve over time.....	45
11. In Conclusion	45
12. Glossary	46
13. References.....	48
14. Appendices	50

1. Executive Summary

Staffordshire Wildlife Trust was commissioned by Cannock Chase District Council to carry out a strategic assessment of the District's biodiversity and habitat networks. This document outlines the existing picture of the District's nature network and describes key locations where habitats may be created or enhanced to contribute to nature's recovery (the Nature Recovery Network), as well as delivering against objectives set out in national planning policy legislation.

Existing data, previous biodiversity opportunity mapping, along with local, regional and national landscape designations and projects were taken into in this assessment methodology.

The methodologies developed aim to deliver against national policies and are used in conjunction with the Department for Environment Food and Rural Affairs biodiversity metrics 2.0 (beta test version) to carry out a strategic broad scale District level spatial assessment of the 'quality components' described in the metric. This included:

1. Habitat distinctiveness
2. Strategic significance (of habitat areas)
3. Habitat connectivity

By using the results above and specific habitat connectivity modelling software it has been possible to define Habitat Connectivity Opportunity (HCO) areas based on habitat types. This is an important next step in identifying areas which possess existing good habitat connectivity and where there is potential for future habitat creation or restoration to contribute to a more successful nature recovery network.

The HCO areas are described in terms of their key opportunities, threats, key species and other habitats which they support along with any potential 'add-on' benefits (e.g. ecosystem services) which could be derived from having well-connected diverse habitat networks contributing to a healthy nature recovery network.

The opportunity map is not static and as physical habitats change on the ground and are subsequently mapped and monitored, the map itself will evolve with these updates. The opportunity areas themselves are where work to enhance habitats can be focussed, where the opportunity to get the greatest benefits lies.

Analysis and opportunity areas mapped within the nature recovery network completed as part of this study are to a fine scale and based around a more robust defensible methodology that can more clearly deliver against National Planning Policy Framework and Planning Policy Guidance objectives, as well as those likely to emerge as outlined in the Environment Bill (House of Commons, 2019).

2. Introduction

Staffordshire Wildlife Trust were commissioned by Cannock Chase District Council to carry out a strategic assessment of the District's biodiversity and habitat networks, to form part of an evidence base in order to ensure biodiversity is an integral part of policy development.

The project encompassed some additional phase one habitat survey; habitat connectivity analysis mapping and Local Nature Recovery mapping. These elements will enable the District to address the requirements articulated within para 170 and para 174 of the National Planning Policy Framework 2019 - *To provide for the protection and support enhancements to the Districts natural environment through the identification, mapping and safeguarding the components and enabling connectivity, interpretation and integration of the natural resources to deliver overall net gain for biodiversity.*

Geographic Information System (GIS) tools were used in the creation of the Nature Recovery Network maps. A GIS software package called MapInfo was used for the digitisation of habitats and network maps. A second piece of software called Condatis was used to model potential species movement through habitat connectivity, the outputs of which were digitised in MapInfo. These technologies are relatively recent and are continuing to develop therefore the results generated by these software are likely to become more sophisticated and accurate over time.

Ecosystem services are the freely gained human benefits provided by natural processes in properly functioning ecosystems for example flood prevention or climate regulation. Whilst ecosystem services have not been assessed as part of this project it is important to recognise that a strong Nature Recovery Network will also provide strengthened Ecosystem Service benefits.

Policy Background

Since previous opportunity mapping for the District was carried out over 10 years ago, there have been considerable changes both in the knowledge, practical assessment and planning of landscape ecology as well as new policy requirements for councils to consider how to protect, enhance and restore biodiversity and the services that it provides.

Key stimuli in updating spatial environmental objectives were documents such as *Making Space for Nature: A review of England's wildlife sites and ecological networks* report by Lawton *et al.* (2010), the Government's 25 Year Environment Plan (2018) and most recently The Environment Bill.

The fundamental principles behind the Making Space For Nature report are for England's ecological network to be 'more, bigger, better and joined' to ensure the survival of species in the face of multiple pressures at a range of scales. The Government's 25-year environment plan puts more impetus on the statutory need to consider the conservation of biodiversity and ensure that it is effectively accounted for through the spatial planning system and the recently published DEFRA Environment Bill.

The emerging Environment Bill sets out environmental principles directed toward the restoration and enhancement of nature and plots a course for how these should be achieved through Nature Recovery Network mapping at a local level ('Local Nature Strategies') and will be a key document in driving the way that these networks are developed and delivered.

Additionally, updated guidance through the National Planning Policy Framework (NPPF) (Ministry of Housing, Communities and Local Government, 2019) and Planning Practice Guidance (PPG) (Ministry of Housing, Communities and Local Government, 2019) have served to put more emphasis on the protection and conservation of nature and our natural resources through spatial planning, providing further justification for the need to have a Nature Recovery Network in place to create a roadmap of where these enhancements could and should go.

The Environment Bill intends to bring in mandatory biodiversity net gain provision, and the use of biodiversity metrics to assess this provision. The Biodiversity Metric 2.0 provides a means of assessing changes in biodiversity value (losses or gains) brought about by development and changes in land use management. The metric is habitat based and gives consideration to improved ecological connectivity.

Habitat opportunity maps are designed to be used in conjunction with biodiversity metrics, in particular Biodiversity Metric 2.0. They can also be used to both inform the metric and target the location and application of future ecological enhancements contributing to a functional nature recovery network.

3. Review of previous biodiversity opportunity mapping assessments

Before using new methods of spatially assessing and targeting opportunities for the enhancement of biodiversity, it is important ensure that they:

1. Can provide additional detail which complements existing objectives.
2. Can provide standalone detail in addition to existing objectives which can be used as evidence in its own right.

The previous biodiversity opportunity mapping assessment carried out for Cannock Chase District (Appendix H) was reviewed and used as a benchmark to compare the results of the updated methods and models used, as part of the new nature recovery network assessment.

The previous methodologies used for biodiversity opportunity mapping throughout the county were based largely on local expert knowledge and stakeholder engagement via practical mapping exercises. Stakeholders and local experts were asked to highlight areas geographically that they saw as priorities for specific habitat and species conservation within a local authority (LA) area. The results of this were sense checked by Staffordshire Wildlife Trust, using available environmental data synthesized into a combined opportunity map and report. These defined spatial landscape areas and detailed conservation priorities within each LA area. The resulting map was effective in that by using expert knowledge, alongside ecological data, as opposed to purely relying on available datasets it was possible to produce an opportunity map with zero white space (areas of a map which have no information). This something which is important to inform decision making on a broad scale and to develop a nature recovery network.

Whilst new methodologies provide a level of additional detail and scientific rigour, local expert knowledge is still vital to ensure that spatial analyses and metrics make sense in a local context.

4. Existing evidence base review

Gathering a robust evidence base is the first step to inform the assessment of opportunities to enhance habitats. The evidence base enables the production and justification of opportunity areas and the assessment of the potential to provide ecosystem services. An inventory of available datasets is one way of bringing together an evidence base forming a platform on which to carry out further analysis.

4.1 Available environmental datasets

A list of relevant Geographic Information Systems (GIS) datasets available for use in completing the mapping assessment are listed in Appendix C, these include datasets held by Staffordshire Ecological Record (SER).

Using the data held by SER and SWT along with publicly available datasets accessible either through an Open Government License (OGL) or through Creative Commons licensing identified in the Natural Capital Committee workbook (Natural Capital Committee, 2017) it was possible to bring together a comprehensive inventory of datasets for review.

Many of the datasets in the inventory are raw or primary data generated directly from information gathered from either desk based or field surveys and remote sensing.

Identification of the coverage and quality of a local authority's environmental dataset provides the baseline for analysis. The results then show how the environment can be protected and enhanced to continue to provide both public and further environmental benefits. By aggregating datasets it is possible to build a composite assessment of the biodiversity within an area without any white space

4.2 Phase 1 habitat mapping via aerial photography interpretation

The District was subject to a Phase 1 habitat mapping assessment were areas where there was no existing Phase 1 habitat data available, i.e. 'white space' as highlighted in the consultant brief.

The overall amount of white space in the District was relatively small, and it was possible to complete the mapping exercise via a desk based methodology. This involved manually assessing the white space against aerial photograph imagery in a GIS package, mapping habitats that can be seen and identified from the aerial photographs. Staff experienced in both on the ground and aerial interpretation habitat survey completed the mapping. It was not possible, due to the time of year and project timeframes to carry out ground-truthing so confidence in the digitised habitat data is subject to limitations (see section 4.5 for full data limitations).

Cannock Chase District has already been extensively habitat mapped using on the ground surveys between 2006 and 2008, this covered a large proportion of the Districts semi-natural habitats leaving only a limited amount of white space remaining to habitat map.

The completion of the current mapping exercise has resulted in an almost complete¹ Phase 1 habitat dataset for the local authority area. This dataset is a composite of habitat data from a wide range of years.

A breakdown of the extent of the Phase 1 habitats dataset can be found in Appendix A

4.3 National Character Areas in Cannock Chase District

There are 159 National Character Areas (NCA) in England, each of which is distinctive with a unique 'sense of place'. These broad divisions of landscape form the basic units of cohesive countryside character, on which strategies for both ecological and landscape issues can be based. The Character Area framework is used to describe and shape objectives for the countryside, its planning and management. These NCA areas are very broad and can encompass a number of different objectives depending on the designated landscape and its respective character, biodiversity and challenges.

Almost all of Cannock Chase District is covered by the Cannock Chase and Cank Wood NCA; a small sliver of land in the north of the District lies within the Needwood and south Derbyshire Claylands NCA, however this is more likely due to the coarseness of the landscape level NCA boundaries rather than any meaningful changes or differences in the overall character or distinctiveness of the area.

The Cannock Chase and Cank Wood NCA identifies 3 broad opportunities for the conservation and enhancement of the areas biodiversity and character. The opportunities identified are:

1. Expand lowland heathland to increase habitat connectivity, improve resilience to climate change and improve water quality.
2. Manage, expand and enhance the network of green infrastructure such as woodlands, restored mining sites, parklands and canal routes, to increase biodiversity, access and recreational use and increase understanding of the area's rich industrial heritage, particularly geodiversity.
3. Conserve and enhance the essential character of this varied landscape, which includes the Cannock Chase Area of Outstanding Natural Beauty, the Forest of Mercia and the urban conurbation of Birmingham and the Black Country, to maintain food and timber production where possible, enhance landscape, sense of place and tranquillity; and increase resilience to climate change.

¹ There may be small gaps in habitat data arising from digitising error or difficulty of creating a seamless fit based on existing data and newly created data. Gaps present in this mapping data are largely a result of difficulties associated with mapping none semi-natural habitats (i.e. man made or artificial surfaces) which are of limited ecological importance.

For each of these three broad level opportunities several smaller sub-objectives have been described to contribute toward the fulfilment of the wider opportunity. These sub-objectives will be used in this report to help define more targeted objectives within the Habitat Connectivity Opportunities area mapping in section 7 to ensure that local objectives and opportunities align with those identified nationally.

In addition to the NCAs are the Staffordshire Ecosystem Action Plan Areas (EAPs) which were created as part of the Staffordshire Biodiversity Action Plan (SBAP). The objectives behind the EAPs are to provide a sustainable living and working environment that benefits both people and nature, by replacing conventional Habitat and Species Action Plans with 14 EAP areas across the county to prioritise conservation management at a landscape level and contribute to national, regional and local conservation targets. The EAPs were used alongside the NCAs to refine the creation of a detailed opportunities map and to make sure that priorities align at both local, county and national level.

4.4 Minerals Safeguarding Zones in Cannock Chase District

There is a need to consider minerals safeguarding zones in the nature recovery network mapping as these areas present both challenges and opportunities from a nature conservation perspective. Whilst the nature recovery network mapping is not spatially aligned on the minerals safeguarding zones. It is important to recognise that that these areas could potentially have a huge impact on the nature recovery network in future, either positively, negatively or both. Where overlaps exist between the maps, there is opportunity to deliver multiple outcomes.

Whilst the likelihood is that much of the safeguarding area will never undergo any mineral extraction, planning any developments within them must be considered to ensure that this will not prevent mineral extraction on potential future extraction sites.

It is possible that high quality habitats may be lost as a result of mineral extraction, a mineral safeguarding zone may also provide protection to important habitats by protecting them from other types of developments.

Post extraction habitat restoration should be guided by the nature recovery network map to create habitats which will most suitably contribute to habitat connectivity within the landscape. In doing this it is possible for mineral extraction sites to contribute to the creation of a diverse and well-connected landscape.

Land within the minerals safeguarding zone may never be actively worked in the long term but could be of great value in terms of contributing to diverse well connected habitats and landscape either if no mineral extraction were to occur or through well planned sympathetic habitat restoration.

4.5 Data used and limitations

It is important to determine the limitations of any datasets identified to ensure that the best possible dataset(s) are used to give the best outcomes for connectivity mapping.

A number of factors can influence whether a dataset is suitable, for example age of the data and whether the data is in a format which can easily and readily be interrogated are crucial in deciding which datasets should be used.

Following a data review the combined habitat map produced during the earlier stages of this was used as a primary baseline from which as this represented the most complete habitat dataset for the area and would easily work with the preferred methodologies to generate the desired technical outputs detailed in sections 5-7.

Several datasets were used in the production of the Nature Recovery Network mapping, justification on their use and relevant limitations can be found in Appendix B.

A full inventory of available datasets has been collated (Appendix C) where each dataset was allocated a 'confidence' rating based on that particular datasets desirability and reliability which helps to justify a hierarchy of use i.e. where there is commonly high desirability and reliability there is a higher 'confidence' in that dataset and it is placed higher in the hierarchy than a dataset which for instance may have a high desirability but a low reliability.

5. Mapping the opportunities to enhance habitats for biodiversity

The first step in analysis to establish opportunities for nature's recovery is to take the data evidence base established previously and carry out a variety of habitat analyses to determine distinctiveness /character for use within other recognised methods (for example, biodiversity metric 2.0 etc...). Furthermore, using the evidence base to apply methods to identify strategic habitat areas and habitat connectivity opportunity areas in relation to creating a robust nature recovery network for the District.

By utilising the knowledge of the county's habitats and species, experience of technical GIS systems and data management, coupled with the available datasets identified in the evidence base, it was possible to produce a number of outputs which are robust, challengeable and can deliver the Districts nature recovery network.

5.1 Habitat distinctiveness mapping

Habitat distinctiveness mapping is one of several elements included within the biodiversity metric 2.0 (Crosher et al. 2019) by using habitat as a proxy for wider biodiversity value via associating and scoring different habitat types according to their relative biodiversity value. An example of this would be irreplaceable ancient woodlands scoring very highly (higher biodiversity value) whereas intensively managed amenity grassland or highly improved agricultural arable land score lower (lower relative biodiversity value).

The criteria used for the creation of the habitat distinctiveness map was based on the Biodiversity Metric 2.0 Beta test (Crosher et al., 2019) which loosely defines what habitats are included within each distinctiveness band. These metrics are currently emerging and form the basis of the Environment Bill (house of Commons 2019), but represent the most comprehensive set of standards for which to base the distinctiveness mapping on.

The distinctiveness map (map 1) was produced using Phase 1 habitat data by associating a distinctiveness value to each specific habitat type (e.g. arable land) in a GIS package based on guidance provided in Crosher et al. 2019, selecting and isolating the habitats spatially into the 5 respective distinctiveness bands. Further ratification to the irreplaceable habitats in the very high distinctiveness band was completed by use of priority habitat inventory (Ancient Woodland Inventory) boundaries. A spatial GIS file was produced for each distinctiveness band.

Further detail of the habitat distinctiveness mapping and the breakdown of habitats included within each distinctiveness band can be found in Appendix D.

Habitat distinctiveness mapping provides multiple uses outside of the biodiversity metric 2.0, including:

1. Identifying areas of high biodiversity value that are a priority for protection and expansion within a local plan whilst working in line with biodiversity mitigation hierarchy (avoid, minimise, remediate, compensate).
2. Flagging areas that may contain medium value (semi-natural) habitat. These could be highlighted in policy as requiring a comprehensive biodiversity evaluation if they are put forward for planning purposes (based on mitigation hierarchy). Biodiversity offsetting/compensation may be required in these areas if they are developed.
3. Identifying possible wildlife corridors which can be highlighted and designated as part of a local plan/Green Infrastructure Strategy. These areas could be the target of restoration projects/funding/aspirational opportunity areas funded through development compensation (obviously the allocation of funds is based on broad scale spatial analysis as opposed to the methods of calculating the offsetting requirement of a specific site).

Planning policy supports application of the mitigation hierarchy which determines a hierarchy of actions when using the biodiversity metric 2.0. This may mean retaining habitats in situ or avoiding habitat damage. It is easier to achieve biodiversity net gains where habitat impacts are avoided due to the way that habitat creation and enhancement risks are accounted for. The mitigation hierarchy is in the desirability order as follows:


- **Avoid** – Where possible habitat damage should be avoided
- **Minimise** – Where possible habitat damage and loss should be minimised
- **Remediate** – Where possible any damaged or lost habitat should be restored
- **Compensate** – As a last resort, damaged or lost habitat should be compensated for

The mitigation hierarchy corresponds with the habitat distinctiveness mapping, e.g. very high distinctiveness habitats such as irreplaceable ancient woodlands should be avoided from development, low and medium distinctiveness habitats could be restored to a higher quality habitat.

The habitat distinctiveness mapping is based on available habitat data and the designated nature conservation site boundaries for the District, including UK Biodiversity Action Plan (UKBAP) and priority habitat areas.

Habitat distinctiveness mapping does not include species explicitly. Instead, it uses broad habitat categories as a proxy for the biodiversity 'value' of the species communities that make up different habitats. The metric does not change existing levels of species protection and the processes linked to protection regimes are outside the scope of the metric.

Habitats are assigned to distinctiveness bands based on an assessment of their distinguishing features including for example rarity (at local, regional, national and international scales), and the degree to which a habitat supports species rarely found in other habitats.



Map 1 Habitat distinctiveness map for Cannock Chase District (2019)

5.2 Habitat distinctiveness mapping limitations

The distinctiveness mapping has been carried out using a desk-based methodology utilising available habitat datasets at a landscape scale with a view of being able to quickly determine on a wider scale the likely impacts of a development. As such the landscape level distinctiveness map in some cases may not provide an accurate account of a sites full habitat distinctiveness at a finer scale (for example at site level). Due to this, developments requiring distinctiveness mapping as part of biodiversity offsetting net gain analysis should be subject to a thorough Preliminary Ecological Assessment (PEA) to determine the full extent of in situ habitats and the expected biodiversity impact of any potential habitat loss or damage.

5.3 Biodiversity metric 2.0²

The DEFRA Biodiversity metric 2.0 is designed to quantify biodiversity to inform and improve planning, design, land management and decision-making.

The metric can be used to both:

- Assess or audit **the biodiversity unit value** of an area of land and
- to **calculate the losses and gains** in biodiversity unit value from changes or actions which affect biodiversity, for example building houses or a change of use in a land holding.

The biodiversity metric 2.0 has 4 'quality components' namely:

- **Distinctiveness** – based on the type of habitat present. For example, modified/amenity grassland is given a score of “2”.
 - Distinctiveness is determined by the habitat distinctiveness mapping (see section 5.2).
- **Condition** – based on the quality of the habitat. This is determined by condition criteria set out in the technical supplement.
 - This cannot be achieved as part of this exercise due to the difficulty of determining condition from a desk based methodology.
- **Strategic Significance** – based on whether the location of the development and or off-site work has been identified locally as significant for nature.
 - Strategic significance is determined by the individual habitat strategic areas and the combined strategic areas map (see section 5.3).
- **Connectivity** – based on the proximity of the habitat patch to similar or related habitats.
 - Connectivity is determined by combined strategic areas map & habitat connectivity opportunity maps (see sections 5.3 & 6).

² The DEFRA Biodiversity metric 2.0 is currently in a beta testing period, the final metric may be different to the metric used in this report.

Through the current study 3 of the 4 quality components have been assessed and defined at a District scale, the only exception being habitat condition which cannot realistically be assessed through a desk based methodology and would require further ground-truthing to determine actual unit values (for example through a Preliminary Ecological Appraisal (PEA)).

5.4 Strategic Habitat Areas

The Strategic habitat area methodology we have applied was developed and is currently being implemented by Warwickshire County Council (WCC) and was developed in partnership with Warwickshire Habitat Biodiversity Audit (WHBA), The University of York and Warwickshire Wildlife Trust. The methodology forms part of WWCs Sub Regional Green Infrastructure Strategy³ and is used in targeting areas for habitat enhancement through biodiversity offsetting compensation.

This methodology was chosen for this mapping assessment because it can be relatively easily applied with the habitat data available; it is robust having been peer reviewed during development, it is already in use by an adjacent local authority and it is based on the fundamental principles of habitat connectivity identified in Lawton et al. (2010).

The mapping works by assessing the proportion of broad habitats e.g. woodland, grassland, heathland etc. within an area to determine whether these are 'strategic', 'semi-strategic' or 'non-strategic' for the creation or restoration of further habitat based on the proportion of habitat already present in the area.

The strategic habitat areas were produced using the composite Phase 1 habitat data identified in the evidence base review. Firstly, specific higher quality habitats were selected and isolated from the composite Phase 1 habitat map (e.g. heathlands or species-rich grassland). The proportion of the selected habitats that overlap individual Ordnance Survey 1km grid squares was then calculated in a GIS package and each square subsequently classified into one of the area bands below, based on the area of habitat overlapping the 1km square. Specific details on the strategic areas are listed in Appendix E.

The strategic habitat areas can be viewed as a hierarchy when it comes to the creation of a particular type of habitat:

1. **Strategic areas** are key areas to focus habitat creation or restoration. There is some high quality semi-natural habitat but additional high quality semi-natural habitat would improve the function of the network.
2. **Semi- strategic areas** are the next preferred areas in terms of habitat creation – These areas already have a relatively large area of high quality semi-natural habitat but more would still be of benefit.
3. **Non-strategic areas** are where there is very little or no high quality semi-natural habitat where it would be difficult to create enough high quality semi-natural habitat

³ <https://www.warwickshire.gov.uk/directory-record/2160/sub-regional-green-infrastructure-strategy>


for it to be functional. (This is not to say that semi-natural habitats should not be created in this area but that it is lower in the overall hierarchy).

The strategic area mapping described will be crucial in delivering the fundamental principles in Lawton et al. (2010).

An overall strategic areas map was produced based on the combination of all the habitats analysed as part of the strategic mapping exercise (map 2). For this map, the criteria for strategic and semi-strategic areas have been swapped so that strategic areas are those with the highest amount of overall habitat. By altering the methodology in this way it is possible to create a coarse overall 'connectivity map' by highlighting the areas with highest combined overall habitat availability and connectivity as opposed to those areas where it is best to create habitats.

The strategic areas are not static and are merely a snapshot in time, changes are an inevitable part of the mapping as available habitat data changes. To an extent the strategic areas mapping is self-fulfilling, as opportunities to enhance habitats described by the map are practically implemented on the ground, mapped through subsequent monitoring and the new habitat data being incorporated into future maps will influence future changes in the areas on the map (described in more detail in section 10.2).

All strategic areas for each of the habitat types assessed are supplied as digital GIS appendices to this report.



Map 2 Combined strategic areas map for Cannock Chase District (2019)

6. Establishing the Habitat Connectivity Opportunity Areas (HCO) for Cannock Chase District

The strategic areas mapping described previously still leaves gaps between areas deemed to be strategic or semi-strategic for a particular habitat type, therefore the creation of habitats solely within these areas may still end up leaving isolated patches habitats which potentially do not link to one another within a landscape. In the interests of driving habitat creation in the direction of connecting these isolated spaces it is important to map an aspirational 'ideal' connected habitat network to work toward: A Nature Recovery Network.

Using local knowledge coupled with additional datasets including soils, nature conservation site boundaries, Staffordshire Biodiversity Action Plan (SBAP) Ecosystem Action Plan Areas (EAPs (Appendix J) and priority habitat inventories along with a piece of ecological modelling software called Condatis (Wallis & Hodgson, 2012), it was possible to further scrutinise and refine the strategic areas map to define comprehensive Habitat Connectivity Opportunity (HCO) areas map for the District based on individual habitats.

The HCO areas add another dimension to the strategic areas modelling detailed previously to define where habitats are both already well connected and equally as crucially broadly identify where to direct the delivery of habitat creation or restoration to create a connected habitat network.

6.1 Habitat Connectivity Opportunity Areas Rationale

The decision to use Condatis to build upon the strategic mapping was in part due to the fact the software has previously been used to identify habitat connectivity in other areas of the county (Churnet Valley Landscape Ecology Pilot Partnership, 2014), where it worked well at identifying rough habitat corridors. Condatis also works on a per habitat basis it is therefore possible to analyse habitat connectivity on an individual habitat basis (A full technical explanation of the Condatis software can be found in Appendix F). Condatis has some limitations in that it only takes into account a single habitat at a time and does not account for other potential connectivity barriers, for example urban areas. It is therefore crucial that these outputs were vetted against other relevant datasets such as soils data; ensuring that identified connectivity opportunities fall in line with the SBAP EAPs areas and that crucially the connectivity opportunity areas correspond with how local expert knowledge would expect the habitat connectivity areas to look in the District, to sense check what is produced by the models.

7. Results

7.1 Habitat Connectivity Opportunity Areas identified

A total of 6 separate Habitat Connectivity Opportunity area types have been identified and mapped covering the entirety of Cannock Chase District:

1. Woodland
2. Grassland
3. Heathland
4. Wetland
5. Urban
6. Pasture and arable land


Each opportunity area is described in terms of its key habitat or habitats. This should not be taken to mean that other habitats are absent from the opportunity area, or that habitats identified as a priority in the opportunity areas should replace existing non-target high quality habitats of a different type.

The Habitat Connectivity Opportunity areas were brought together to produce a combined HCO map for the District (map 3).

For such a small and largely urbanised Local Planning Authority area, Cannock Chase District retains a proportionally very high level of biodiversity within the remaining areas of semi-natural vegetation, particularly in regards to Heathland and Woodland habitats as well as possessing the only designated Area of Outstanding Natural Beauty (AONB) in the county. The AONB is largely undeveloped, with only a handful of small settlements and encompasses most of the District's more semi-natural landscape.

The proximity of the AONB to the urbanised areas of the District presents multiple opportunities and threats in terms of a nature recovery network. Whilst there is clearly a high level of human pressure exerted on any habitats present through recreational activities this also provides a huge opportunity to engage with people both locally and further afield the importance of nature conservation and how a healthy nature recovery network can not only improve wildlife's chances of survival but can also greatly benefit the health and wellbeing of the public.

Each opportunity area is described in more detail in the following sections, along with relevant associated land uses, environmental issues, and the overarching objectives and opportunities for each zone.



Map 3 Combined habitat connectivity opportunity areas map for Cannock Chase District (2019) NB: some of the HCO areas overlap one another which can lead to the colouring of the map being distorted.

7.2 Woodland Opportunity Area

Whilst woodlands in the District on the whole may not necessarily be of semi-natural ancient status, overall coverage of woodland area is very high, the majority of the woodlands in the District are mixed and coniferous plantation woodlands managed for timber production in and around the Cannock Chase AONB on the freely draining glacial sands and gravel soils.

There are 3 sites in the District on Natural England's Ancient Woodland Inventory (AWI) having been continually wooded for the past 400 years, however 2 of these (Chetwynds Coppice and Brereton Hayes Wood, both near Rugeley) are for the most part designated as ancient replanted woodlands meaning they have been replanted at some point within this timeframe. Courtbanks Covert near Prospect Village is the only other site in the District with an AWI designation, all of these AWI woodlands are within the AONB.

The industrial heritage of the District has led to the formation of several areas of woodland where former quarries and collieries since abandoned have undergone natural successional regeneration have since been colonised by scrub and young woodland, mostly dominated by Birch and Willow species along with species favouring more free draining and acid conditions such as Gorses and Broom.

7.2.1 Key Habitats

- Woodlands (e.g. semi-natural ancient woodlands, coniferous plantation)
- Wood Pasture and Parkland
- Dense Continuous and Scattered Scrub
- Lowland Heathland
- Traditional Orchards
- Lowland Meadows

7.2.2 Key species

- The Districts woodlands are important for a number of bird species, both breeding and visitors, these are too numerous to mention individually but a select few have been highlighted below, however this list is not exhaustive.
- Nightjar
- Crossbill
- Cuckoo
- Owl Species
- Bluebell
- Amphibian species (Great Crested Newt, Common Frog, Common Toad), specifically as terrestrial refuge sites.
- Red and Fallow Deer
- Hybrid Bilberry

7.2.3 Threats

- Loss and fragmentation of irreplaceable woodland habitats (ancient woodland inventory sites).
- Both residential and industrial development.
- Inappropriate management of species-rich and/or ancient woodland sites either directly within or surrounding

7.2.4 Opportunities

- Protection of existing sites, particularly ancient woodland inventory sites and woodlands which are designated as Local Wildlife Sites. Planting of further future woodlands on sites which do not already support a priority habitat to improve connections of existing areas of high quality woodland and increase the area of woodlands which are ecologically functional for the species that they support.

<p>these sites leading to deterioration and lowering overall diversity.</p> <ul style="list-style-type: none"> • Loss or deterioration of hedgerows and other associated habitats severing connectivity between woodlands and to other habitats. • Unsympathetic or poorly thought out woodland planting and creation on sites which already support another important habitat of conservation concern, e.g. species rich lowland meadows, causing irreversible loss. • Replanting of ancient woodland sites with species which are not characteristic or native to the area. • Scrub removal either through intensive management regimes or development. 	<ul style="list-style-type: none"> • Encourage relaxed management on the fringes of woodlands to provide a softer edge (e.g. scrub formation) habitat which is able to support both more and a wider diversity of species, particularly birds and butterflies. • Expand the area of existing woodlands. Create new areas of woodland that are in strategic locations and are of suitable size to act as stepping stones between existing woodlands. Woodland expansion and creation must not be detrimental to other high quality habitats for instance diverse grassland habitats. • Use historical maps and data to determine the past extent of woodland areas, particularly where there may still be a rich ground flora in the seedbank for the restoration and expansion of ancient woodland sites. • Planting new and maintaining existing hedgerows to better connect smaller isolated woodlands benefiting species migration and chances of breeding. • Avoidance from or incorporating key woodlands into development sites, this is achievable through mitigation hierarchy in the biodiversity offsetting system. • Restoration of Planted Ancient Woodland sites (PAWS) to native broadleaf or diversification of coniferous woodlands to include more native planting. • Ensure that there is no loss or damage to known wood-pasture or parkland sites. • Identification of, and promotion of the importance of veteran trees, both in woodland and in the wider landscape.
7.2.5 Specific opportunities	
<ul style="list-style-type: none"> • Restoration of the coniferous plantation woodlands around the southern areas Cannock Chase AONB to heathland and/or broadleaf woodland. • Ensure that designated Ancient Woodland Inventory sites at Chetwynds Coppice, 	


Brereton Hayes Wood and Courtbanks Covert are all protected from damage or loss.

- Seek creation and extension of existing woodlands east of Heath Hayes park and the area of land between the M6 and A5 on the edge of the district north of Great Wyrley. This would provide added connectivity opportunity in the more developed areas as well as additional screening.

7.2.6 Opportunities to enhance other benefits

- Flood risk mitigation
- Carbon storage
- Recreation and aesthetic
- Cultural heritage
- Wood fuel, timber and fibre
- Foraging / wild food

7.2.7 Map of Woodland Opportunity Area



Species rich grasslands occur throughout the District and can largely be divided into two categories; Lowland acid grasslands which in general occur as a mosaic with Lowland Heathland and Neutral Grasslands, managed as Lowland Meadow and species rich pastures. Marshy grassland, Floodplain Grazing Marsh and Purple Moor Grass and Rush Pastures do occur in the District, however these are generally restricted to the floodplains of the River Trent in the north of the District, the Saredon Brook in the south of the District and a very limited amount of wet grassland on the periphery of Chasewater.

Lowland Heath Acid Grassland mosaics are most prevalent throughout the AONB area, particularly around Brindley Heath, Brindley Valley and Penkridge Bank. Other areas which possess Acid Grassland/Lowland Heath mosaics are Hednesford Hills, some areas around Heath Hayes, Prospect Village, Wimblebury and Rawnsley.

Semi-natural neutral/mesotrophic grasslands occur throughout the District as a whole, both within and outside of the AONB area, concentrations of which exist around Cannock Wood, Hazelslade, Heath Hayes and Great Wyrley/Norton Canes.

Species rich grassland sites are fewer and smaller in the south-west of the District due to the fact that this area has been subject to the most development, however several species rich sites do survive amongst the urban landscape, in some areas allowing mosaics of habitat to proliferate.

7.2.1 Key Habitats

- Lowland meadows
- Semi-improved Pastures
- Hedgerows
- Arable land
- Open mosaic habitat on previously developed land
- Ponds
- Traditional Orchards
- Coastal Floodplain Grazing Marsh
- Purple Moor Grass and Rush Pastures

7.3.2 Key Species

- Skylark
- Farmland birds
- Bats (specifically Brown Long-eared, Noctule and Pipistrelle species)
- Butterfly species such as Dingy Skipper and Small Pearl-bordered Fritillary.
- Reptiles, specifically Common Lizard and Adder
- Red and Fallow Deer

7.3.3 Threats

- Development pressure
- Poor management of key diverse sites including:
 - Over-grazing
 - Poaching
 - Neglect of Hedgerows
 - Over-cutting of Hedgerows
- Nutrient intensification both from agricultural practices as well as diffuse pollution sources - nitrogen deposition.
- Agricultural intensification

7.3.4 Opportunities

- Ensure that all high quality grassland sites remain in positive conservation management, securing vital areas which can be used as sources of biodiversity into the future.
- Protection of existing high quality grasslands and buffering these from potentially detrimental neighbouring land uses such as intensive farming practices. This could be achieved through encouraged uptake of agri-environment schemes, landowner liaison/education

<ul style="list-style-type: none"> ● Neglect of sites and lack of appropriate conservation management. This is a major threat to diverse grassland sites not only in the District but in the county as a whole. ● Global and local climate change. ● Habitat loss and fragmentation 	<ul style="list-style-type: none"> ● Enhancement of any existing grassland sites or restoration of degraded sites so that they may achieve Local Wildlife Site Status and ensure that the management of these sites persists to ensure that they remain diverse. ● Reversion of arable land to diverse grassland where soils dictate. This is usually only carried out in certain circumstances due to the difficulty and cost associated however there are examples of this being successfully carried out in the District. ● It is critical that areas of high quality grassland are linked with mosaics of other high quality grassland to ensure that species reliant upon these habitats are able to move freely between them. ● Use of Light Detection And Ranging (LiDAR) imagery to identify historical field patterns and features i.e. ridge and furrow to indicate where grassland restoration may be most successful as these areas have not or are unlikely to have undergone any serious agricultural improvement in the past.
----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------


7.3.5 Specific Opportunities

- Protection and enhancement of pockets of more diverse grassland in and around the urban environment, particularly around Green Heath, Hawksmoor, Littleworth, Hawk's Green and Wimblebury.
- Protect, Enhance and restore the areas of grassland heathland mosaic to the east of the district around Cannock Wood as these provide important habitat connectivity on the periphery of the district boundary, particularly to sites such as Gentleshaw Common.
- Restore and enhance grasslands in and on the outskirts of Rugeley to improve species permeability from the larger expanses of semi-natural habitat in the Cannock Chase AONB to the West.
- Utilise suitable designated green space throughout Cannock and Rugeley for the creation or enhancement of species rich grassland. This will provide vitally important connectivity throughout what is otherwise a hindrance for species movement throughout the urban environment.

7.3.6 Opportunities to enhance other benefits

- Pollination
- Recreation and aesthetic
- Cultural heritage

7.3.7 Map of Grassland Opportunity Area



Cannock Chase District is a stronghold for Lowland Heathland in Staffordshire and acts as a core connecting site between the northern extent of the Cannock Chase AONB in Stafford Borough which has a large core area of Lowland Heath habitat and areas of heathland further south in Lichfield District such as Gentleshaw and Chasewater through to the Greater Birmingham and Black Country area and the heathlands of Pelsall Common and Sutton Park on the county boundary.

The areas of heathland through Cannock Chase District act as a backbone of connectivity providing 'stepping stones' to allow species associated with this habitat to move through the landscape.

Significant areas of heathland are present both within and outside the AONB area, often occurring as a mosaic with other high quality habitats such as Lowland Acid Grassland, scrub and brownfield habitats known as "Open Mosaics on Previously Developed Land" (OMPDL) which are also an important feature of the District. Specific areas of note for the presence of this habitat are Brindley Heath, Brindley Valley, Penkrige Bank, Rawnsley Hills, Hednesford Hills and Heath Hayes. The area of heathland supported within the District provides fulfils the criteria of a number of species with specific habitat requirements.

7.4.1 Key Habitats

- Lowland Heathland
- Woodlands
- Arable (buffer strips, set-aside etc.)
- Lowland Acid Grassland
- Lowland Meadows
- Open mosaic habitat on previously developed land
- Scrub
- Ponds

7.4.2 Key Species

- Heather
- Nightjar
- Fly agaric
- Small Pearl-bordered Fritillary
- Green Hairstreak Butterfly
- Reptiles and Amphibians
- Bog Bush-cricket
- Welsh Clearwing Moth
- Hybrid Bilberry
- Aculeate Hymenoptera (specifically solitary bees and wasps)

7.4.3 Threats

- Pollution both from acute and diffuse sources leading to nutrient intensification – e.g. Nitrogen loading.
- Lack of management, improper management or neglect leading to scrub encroachment.
- Tourism and recreational pressure.
- Mineral extraction.
- Agricultural intensification, both on and surrounding core areas of heathland.

7.4.4 Opportunities

- Protection of existing areas of high quality Lowland Heath through sympathetic management and ensuring that positive management continues and prevent degradation due to neglect.
- Seek to create areas of new Heathland in key sites. This could be through development sites as part of biodiversity offsetting mitigation, reverting plantation woodland stands into areas of heathland post harvesting similar to those carried out by the

<ul style="list-style-type: none"> • Urban development. 	<p>Connecting Cannock Chase project or through incorporating into existing habitat management such as arable field margins or relaxing the management regime in pastures etc.</p> <ul style="list-style-type: none"> • Regeneration of former sand and gravel sites by inoculation with heather seed and brash to kick start habitat formation and secure sympathetic management of these sites in future. • Mitigate potential impacts of recreation pressure on sites such as Cannock Chase AONB and Gentleshaw Common to ensure that the habitats and species which exist there can thrive, but can also be enjoyed by those who live in and visit the area. • Ensure that sites listed on Natural England's Heathland Inventory are appropriately managed and monitored to retain key connectivity sites within the District.
------------------------------------------------------------------------	--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------


7.4.5 Specific opportunities

- Ensure heathland sites within the Cannock Chase AONB area are protected, and seek to expand these areas of heathland/acid grassland mosaic through habitat creation and restoration.
- Ensure the heathlands at Hednesford Hills are protected and well managed, this is a key stepping stone site for this habitat in the district, connecting the heathlands further north in the boundary of the Cannock Chase SAC/AONB and those further east and south around Chasewater and Gentleshaw.
- The land to the east of Heath Hayes park/Wimblebury and south of Prospect Village is a priority for heathland creation and restoration acting as a key corridor in the link between Cannock Chase to Sutton Park via the areas of heathland at Chasewater and Gentleshaw Common. The area of land at Bleak House will also be a suitable site for the creation of wet heath habitats benefitting connectivity of similar habitat between Gentleshaw, Chasewater and Sutton Park.
- Continuation of objectives laid out in the connecting Cannock Chase project and use these areas as a positive example to establishing conservation grazing on suitable sites to ensure that the sensitive heathland habitat is in favourable condition and remains so.

7.4.6 Opportunities to enhance other benefits

- Pollination
- Cultural Heritage
- Carbon Storage
- Flood risk mitigation
- Recreation and Aesthetic

7.4.7 Map of Heathland Opportunity Area



7.5 Wetland Opportunity Area

Wetlands are limited in the District however there are a number of watercourses and wetland sites which support both local and national priority species. Only one major river flows through the District, a short 2km stretch of the River Trent to the north which acts as the District boundary between Cannock Chase and Lichfield Districts.

Most wetland areas in the District are small, confined to small brook courses and streams as well as small ponds and pools, Chasewater reservoir lies just over the District boundary in Lichfield District to the south-east and has a range of diverse wetland habitats.

Notable areas of importance are the headwaters of the Upper Sherbrooks, Shropshire, Rising, Old, Stony and Stafford Brooks, and Hednesford Brickworks all of which support populations of protected species important at both local and county level.

Cannock Extension canal SSSI in the south of the District is important for a number of aquatic plant species including protected species and those of national and local importance.

7.5.1 Key Habitats

- Woodland
- Grassland
- Pasture
- Arable
- Urban fabric/mosaic habitats

7.5.2 Key Species

- Otter
- Great Crested Newt
- Freshwater White-clawed Crayfish
- Harvest Mouse
- Reptiles and Amphibians
- Floating Water-plantain
- Red-eyed damselfly
- Emperor Dragonfly

7.5.3 Threats

- Mineral extraction.
- Pollution from acute and diffuse sources.
- Poor land management, livestock in and near watercourses and waterbodies, soil erosion leading to eutrophication of water bodies and loss of habitat in watercourses.
- Historic deepening and straightening of watercourses, meaning that rivers and streams lack natural features such as gravel beds. Water is disconnected from floodplains.
- In some areas removal of tree cover and grazing leading to habitat degradation.

7.5.4 Opportunities

- Protection of existing high quality wetland sites particularly those with a nature conservation designation. This will be achieved through the identification of environmental issues for example pollution from agricultural run-off and subsequent remediation for instance through Rural SuDS. These sites should be buffered from any potential sources of damage both through creation of habitat around key sites to provide a 'soft edge' habitat and landowner liaison to address issues.
- Identification of the most suitable locations for the targeting and prioritisation of further wetland creation and enhancements in connection with implementation of SuDS schemes and

<ul style="list-style-type: none"> ● Lack of understanding of the need to protect water throughout the catchment including areas where there are no obvious watercourses. ● Global and local climate change. ● Loss of 'coarse' habitat to development or agricultural intensification which would otherwise impede the flow of water leading to: <ul style="list-style-type: none"> ○ Increased flood risk. ● Invasive Non Native Species (Himalayan Balsam (<i>Impatiens glandulifera</i>), Parrot's-feather (<i>Myriophyllum aquaticum</i>), <i>Azolla</i> sp, <i>Crassula helmsii</i> etc.). ● Disease/Biosecurity. 	<p>natural flood management.</p> <ul style="list-style-type: none"> ● Seek opportunities to deliver natural flood management to address flood risk as well as provide additional areas for habitat provision. ● Look for opportunities to carry out river reprofiling/naturalisation, improve flood storage and provide additional habitats suitable for a range of species particularly breeding waders and wintering wildfowl. ● Use historical maps and LiDAR information to identify historical wetland and river features, sluices, water meadows etc. which could potentially be restored to deliver both flood risk mitigation and habitat improvements. ● Use flood models to dictate where work can be targeted to both deliver improved flood mitigation as well as deliver further habitat works ● Ensure that sand and gravel quarry extraction sites are effectively restored and provide additional benefits for wildlife. There is a huge opportunity here to deliver biodiversity and flood mitigation objectives. ● Effective mitigation for the loss Great Crested Newt (GCN) habitat as a result of development. (Priority areas for the creation of compensatory pond clusters would need to be addressed at a finer scale using Great Crested Newt (GCN) metapopulation data and modelling).
--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

7.5.5 Specific opportunities

- Ensure that habitats on the Rising Brook and Headwaters of the Shropshire Brook are protected and enhanced to protect important populations of native White-clawed Crayfish. Carry out regular monitoring of populations and ensure that thorough biosecurity measures are in place to prevent disease transmission.
- Improve education, awareness and interpretation of biosecurity risk, particularly in the Cannock Chase AONB especially in relation to transmission of Phytophthora which infects Bilberry and crayfish plague which can have drastic implications on White-clawed Crayfish populations.
- Protection of habitats supporting nationally important plant populations on the Cannock Extension Canal in the south of the District. Buffer the site through the creation of beneficial habitats which will both help to protect the site from detrimental


external factors and help provide additional connecting habitat. Encourage sympathetic management of surrounding land to prevent run-off leading to either pollution of the site or nutrient enrichment which could lead to a loss of diversity.

- Retain, improve and expand wetland habitats at Hednesford old brickworks for the benefit of Great Crested Newt populations.
- Seek to implement Natural Flood Management (NFM) techniques in the headwaters of the Shropshire Brook, Rising Brook to slow the flow of water and alleviate any potential flooding issues further downstream. By using NFM techniques this should also provide additional opportunities for the creation of habitats beneficial to a range of species as detailed above.
- Enhancement and creation of wetland areas upstream of Chasewater reservoir (this may include habitats such as wet heath), particularly around Bleak House. This should improve habitat connectivity of wetlands in the area between Gentleshaw Common and Chasewater Reservoir.

7.5.6 Opportunities to enhance other benefits

- Flood risk mitigation
- Water quality
- Recreation and aesthetic
- Cultural Heritage

7.5.7 Map of Wetland Opportunity Area



7.6 Pasture and Arable Opportunity Area

Pastures and Arable land are relatively limited in the District, mostly present in the far north and far south with some holdings interspersed throughout the remainder of the District, mostly in the east. Arable production only accounts for a small proportion of the Districts land cover (approximately 9%). Large homogenous blocks of intensive arable and pasture are rare in the District, generally intensive farmland is interspersed amongst areas of urban development and semi-natural vegetation or plantation forestry.

7.6.1 Key Habitats

- Grassland
- Woodland
- Hedgerows
- Mature and veteran trees

7.6.2 Key Species

- Barn Owl
- Brown Hare
- Harvest Mouse
- Polecat
- Grey Partridge
- Wall Brown butterfly

7.6.3 Threats

- Habitat fragmentation.
- Agricultural intensification.
- Urban encroachment.
- Pollution of waterways.
- Loss and deterioration of ponds for example through changes in water management or nutrient intensification.
- Improper management e.g.
 - Over-grazing
 - Poaching
 - Neglect of hedgerows
 - Over-cutting of hedgerows

7.6.4 Opportunities

- There are a wide range of opportunities for more intensively farmed agricultural land ranging from very small interventions such as leaving one corner of an arable field as set aside to provide feeding opportunity for farmland seed eating birds to large whole farm scale interventions for example reversion of large areas of arable land into diverse grassland. Obviously the scale of the intervention is down to what is practical and ultimately what is desirable, cost effective and sustainable in the eyes of landowners and land managers.
- Link up existing semi-natural habitats through the creation of habitat corridors and networks using hedgerows, arable field margins and watercourses where possible.
- Reversion of arable to other habitats with a higher biodiversity value for example species rich grassland.
- Encourage uptake or movement toward organic production methods or holistic grazing management over reliance on supplementary feeding or indoor systems for example.

	<ul style="list-style-type: none"> • Where developments are likely to impact on large areas of intensive farmland, ensure that as a result some of the developed area is dedicated to the provision of high quality semi-natural habitats which may greatly improve habitat availability and connectivity within the landscape.
--	------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------


7.6.5 Specific opportunities

<ul style="list-style-type: none"> • Only a small area of the district falls within a pasture and arable connectivity opportunity area on the outskirts of Rugeley, in this area opportunities which are likely to be of most benefit to fit in line with existing land use would be small scale interventions such as hedgerow creation, wider arable margins and changes to land management practices. In future this could possibly be achieved via updated environmental land management schemes through payment for managing land for the benefit of public goods and services.

7.6.6 Opportunities to enhance other benefits

<ul style="list-style-type: none"> • Water quality • Cultural heritage • Food production

7.6.7 Map of Pasture and Arable Opportunity Area



A proportionally large area of the District is urbanised, the town of Rugeley in the north of the District and the conurbation of Cannock, Hednesford, Bridgtown and Norton Canes in the south east of the District as well as smaller settlements such as Cannock Wood and Slitting Mill.

Aside from the town centres of Rugeley and Cannock, the urban area is on the whole discontinuous with good access to high diversity green space a lot of which achieves Local Wildlife Site status. There is a mixture of habitats throughout the urban areas of the District, for example lowland heathland, plantation woodland, semi-natural grasslands and ponds/pools along with the more formal areas of green space with amenity grasslands used for sports and associated recreational activities but are of a more limited biodiversity value.

Having high diversity green spaces already present in the urban areas allows for increased habitat connectivity both between rural and urban areas, improves people's access to nature.

7.7.1 Key Habitats

- Grassland
- Woodland
- Open Mosaic Habitat on Previously Developed Land (brownfield habitats)
- Wetland
- Rivers and streams
- Street trees (particularly those in environments where other green space is lacking)

7.7.2 Key Species

- Hedgehog
- Great Crested Newt
- Slow Worm
- Invertebrates and pollinators
- House Sparrow
- Toads and other amphibians
- Finch species.

7.7.3 Threats

- Habitat fragmentation through the loss of both sources of biodiversity as well as habitat 'stepping stones' and linear pathways which species require to be able to disperse.
- Pollution both from acute and diffuse sources leading to the loss of diversity in waterways etc.
- Urban expansion
- Redevelopment of Open Mosaic Habitats on Previously Developed Land (OMPDL) which are often important sites for a number of species in urban areas.
- Intensive management of urban green spaces leading to:

7.7.4 Opportunities

- The key objectives in these areas is not to connect urban areas together but to enable permeability between rural and urban landscapes, especially where high quality semi-natural habitats exist in close proximity to or within these areas. In doing this it is possible to benefit habitat connectivity but also bring wildlife closer to people.
- Ecological enhancement of existing urban green spaces, for example through improving the diversity of amenity grassland in parks by seed sowing and green hay strewing, enhancement or creation of wetlands in Sustainable Drainage Systems (SuDS).
- Creation of new habitats when

<ul style="list-style-type: none"> ● Invasive species ● Increased flood risk due to increased area of hard impermeable surfaces. 	<p>planning new urban developments, make new developments as green as possible to bring high quality habitats and improve habitat connections in the urban environment. This may include for example green roofs/green walls, wildlife friendly SuDS which can be planted with native wetland species, rain gardens to slow the flow of water.</p> <ul style="list-style-type: none"> ● Ensure that urban green spaces are managed appropriately to provide the best benefits for wildlife and people - this may include relaxing mowing regimes to create and maintain more diverse grasslands, thinning of plantation woodlands to improve structural diversity or invasive species control. ● Ensure that linear features such as canals, old railway lines, road verges, hedgerows are managed for the good of wildlife as these are often critical pathways for biodiversity in and out of the urban environment. ● Provide suitable opportunities in existing and new developments for protected and Biodiversity Action Plan (BAP) species for example bats, hedgehogs and pollinators.
----------------------------------------------------------------------------------------------------------------------------------------------------------	---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

7.7.5 Specific opportunities


- No opportunities identified

7.7.6 Opportunities to enhance other benefits

- Recreation and aesthetic - improved access to and increased number of natural resources.
- Health and wellbeing - improved access to an increased number of natural resources.
- Flood risk mitigation - More green areas lead to increased habitat coarseness which slows the flow of water, Sustainable Drainage Systems (SuDS) schemes increase habitat and hold water away from vulnerable areas.
- ‘Pocket Parks’ encouraging local people to take up management of small urban green spaces to benefit both wildlife and those which live nearby. By adopting multiple pocket parks it is possible to create a stepping stone network throughout the urban environment.
- Urban cooling – suitable tree planting, increased green space and green developments, green walls, green roofs etc.
- Cultural heritage - access to nature and traditional landscapes.

- Public engagement - opportunity to educate people on ecology and the natural world and what people can do to provide space for wildlife in gardens, allotments, local parks etc.

7.7.7 Map of Urban Fabric Opportunity Area




8. Cross boundary habitat connectivity

As habitats and wildlife do not adhere to political boundaries it is important to take into account habitats which exist on the other side of political boundaries to ensure that there is no 'hard edge' where for example a Habitat Connectivity Opportunity area ceases to exist at the edge of a county or District boundary despite there being suitable habitat

Map 4 illustrates this, showing the Habitat Connectivity Opportunity areas combined map including a 2km radius buffer around the District boundary. Despite the buffered radius falling outside of the District and county boundaries habitat connectivity into these areas has been considered as part of the mapping to ensure this 'hard edge' has been avoided. It must be noted however that the HCO areas do not extend large distances into neighbouring authority areas with the ultimate goal that all authority areas will have a mapped Nature Recovery Network which dovetails with this NRN mapping.

The cross boundary HCO areas in neighbouring local authorities may be subject to change based on any future NRN mapping which may be commissioned by the respective local authority in its jurisdiction. At this stage Habitat Connectivity Opportunity areas identified outside of the District should only be considered potential and may be subject to future changes. These areas have been included in this assessment to demonstrate the duty to cooperate across boundaries has been considered in this mapping exercise.



Map 4 Combined habitat connectivity opportunity areas map for Cannock Chase District including a 2 kilometre buffered radius of the District boundary (2019). NB: some of the HCO areas overlap one another which can lead to the colouring of the map being distorted.

9. Practical Application of the maps

The HCO maps detailed are designed to be used in conjunction with the biodiversity metric 2.0, however the habitat connectivity opportunity areas and the bottleneck analysis can be used to both inform the metrics and target the location and application of future ecological enhancements contributing to a functional nature recovery network.

The HCO areas are based around the principle of habitats being ecologically functional and well connected to one another within the landscape. This means that habitats are able to both support a high population and diversity of species, meaning these species have the ability to be able to move freely within the landscape, as a result of good habitat connectivity.

These areas promote the conservation, restoration and enhancement of certain priority habitats, ecological networks and contribute to the protection and recovery of associated priority species within defined geographic areas.

The habitat connectivity opportunity areas mapping has no white space as there are always opportunities for the delivery of habitat creation or enhancement anywhere in the landscape irrespective of whether it has been identified as a connectivity area for a priority habitat or not. Taking this approach ensures that the landscape as a whole can remain permeable for our flora and fauna and resistant to both local and global impacts For full technical details on the principles of HCOs and mechanisms for delivery see Appendix G.

The habitat connectivity opportunity areas identify the key areas where the creation of new habitat is best prioritised to benefit habitat connectivity within the landscape. Targeting additional habitat creation in this way will have the greatest impact on both availability and connectivity of habitat within the landscape as it builds upon areas which already possess some good quality habitats but by increasing their size, quality, coverage and connectivity within the landscape will enable those habitats to become more functional.

Within these areas there are further opportunities to deliver environmental outcomes within existing spatially defined partnership schemes, specifically:

- Cannock Chase AONB
- Cannock Chase Special Area of Conservation (SAC) Partnership

The way that the opportunity areas are generated means that habitat opportunities are not mutually exclusive of one another i.e. there can be overlapping areas for multiple habitat types; for instance an area defined as an opportunity for woodland enhancement may also provide a good opportunity for improving grassland and wetland habitat enhancement and connectivity. The on-site prioritisation of what habitat to create where must therefore rely upon both the opportunity areas as well as local ecological expert knowledge so as not to risk either damaging connectivity or destroying existing good quality habitats.

10. Next Steps

10.1 Habitat connectivity bottlenecks

Bottlenecks highlight the areas of habitat which have the highest 'strain' in terms of supporting connectivity within the nature recovery network. These areas are where there is a high flow of species through an area with relatively few links and over a long distance (i.e. a very concentrated flow of species movement squeezing through a very small area of habitat and being forced to jump large distances between patches of suitable habitat). Producing detailed guidance in how this can best be addressed will enable informed focused positive impacts that directly reduce strain on the habitat connectivity network.

Bottlenecks can be used to determine the optimal locations to create and restore habitats to benefit connectivity and reduce 'strain' on the habitat network. Creating, enhancing and restoring habitat in these locations will not only benefit by reducing strain on the network but also reduces the risk or likelihood of losing what may be an important link in a habitat connectivity network which is already under pressure.

10.2 How the strategic mapping will evolve over time

As previously discussed, the opportunity map is not static and as physical habitats change on the ground and are subsequently mapped and monitored the map itself will evolve with these updates (Appendix K).

11. In Conclusion

The analysis and opportunity areas mapped within the new nature recovery network are much more fine scale and are based around a more robust defensible methodology that can more clearly deliver against NPPF and PPG objectives, as well as those likely to emerge as outlined in the Environment Bill (House of Commons, 2019).

12. Glossary

Term	Definition
	A biodiversity action plan (BAP) is an internationally recognized program addressing threatened species and habitats and is designed to protect and restore biological systems. The original impetus for these plans derives from the 1992 Convention on Biological Diversity (CBD). The UK Biodiversity Action Plan (UK BAP) was published in 1994, and was the UK Government's response to the Convention
Geographic Information System (GIS)	A computer system for capturing, storing, checking, and displaying data related to positions on Earth's surface. By relating seemingly unrelated data, GIS can help individuals and organizations better understand spatial patterns and relationships.
Light Detection And Ranging (LiDAR) imagery	Remote sensing method that uses light in the form of a pulsed laser to measure ranges (variable distances) to the
Local Wildlife Site (LWS)	Local Wildlife Sites are areas with locally significant nature conservation value. They come in all shapes and sizes, from small wildflower meadows and secluded ponds to ancient woodlands. Most are owned by private individuals.
Natural capital	Natural capital can be defined as the world's stocks of natural assets which include geology, soil, air, water and all
Nature conservation site	This is a blanket term is used to describe all sites which have a land use designation relevant to nature conservation or are managed in the interests of nature conservation and wildlife for example, Local Wildlife Sites, SSSI or Nature reserves.
Non-statutory nature conservation site	Non-statutory sites (specifically LWS) receive some protection from development via local planning documents which recognise the need to protect and enhance designated sites and those of interest without a statutory
Site of Special Scientific Interest (SSSI)	Sites of Special Scientific Interest are areas of very high nature conservation value which are legally protected nationally, these sites are normally the best remaining examples of natural habitats and may also have an international designation e.g. Special Area of Conservation (SAC).
Statutory nature conservation site	A site with a designation which is upheld and protected by law e.g. SSSI or SAC
Sustainable Drainage Systems (SuDS)	Sustainable drainage systems (SuDS) are a technical solution to addressing issues that arise with the increasing problem of excess surface water. Originally used in urban areas, they are now used for some roads and towns in rural areas.
	SuDS are always site specific, and require bespoke design that take into account the underlying hydrology, functional

	purposes of the area, and the present and future needs of people using the area.
White space	Areas of a map which have no information, i.e. gaps in a dataset.

13. References

Burkhardt, B., Kandziora, M., Hou, Y. & Muller, F. (2014) Ecosystem Service Potentials, Flows and Demands - Concepts for Spatial Localisation, Indication and Quantification. *Landscape Online*. **34**, pp.1-32.

Churnet Valley Landscape Ecology Pilot Partnership. (2014) *Churnet Valley Landscape Ecology Pilot: Habitat Networks Modelling*. Staffordshire.

Ian Crosher, Susannah Gold, Max Heaver, Matt Heydon, Lauren Moore, Stephen Panks, Sarah Scott, Dave Stone & Nick White. 2019. The Biodiversity Metric 2.0: auditing and accounting for biodiversity value. User guide (Beta Version, July 2019). Natural England

Hölzinger, O. and Everard, M. (2014) *Staffordshire Ecosystem Assessment*. Staffordshire County Council, Stafford.

House of Commons. (2019) *Environment Bill (ordered by The House of Commons to be printed on 15th October 2019)*. [online] Available at: <https://publications.parliament.uk/pa/bills/cbill/2019-2020/0003/20003.pdf>

Lawton, J.H., Brotherton, P.N.M., Brown, V.K., Elphick, C., Fitter, A.H., Forshaw, J., Haddow, R.W., Hilborne, S., Leafe, R.N., Mace, G.M., Southgate, M.P., Sutherland, W.J., Tew, T.E., Varley, J., & Wynne, G.R. (2010) Making Space for Nature: a review of England's wildlife sites and ecological network. Report to Defra.

Ministry of Housing, Communities and Local Government. (2019) *National Planning Policy Framework*. Available at: https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/810197/NPPF_Feb_2019_revised.pdf [Accessed 24 September 2019].

Ministry of Housing, Communities and Local Government. (2019) Collection: *Planning practice guidance*. [online] Available at: <https://www.gov.uk/government/collections/planning-practice-guidance> [Accessed 24 September 2019]

Natural Capital Committee. (2017) How to do it: a natural capital workbook. Version 1. [online] Available at: https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/608852/ncc-natural-capital-workbook.pdf [Accessed 10 June 2019].

Noake, B. Liu, V., Lawley S. & Glaisher, A. (eds) (2016) *The State of Staffordshire's Nature - Summary Report*. Staffordshire Wildlife Trust and Staffordshire Ecological Record. Wolseley Bridge, Stafford.

The Wildlife Trusts. (2018) *Towards a Wilder Britain, Creating a Nature Recovery Network to bring back wildlife to every neighbourhood*. A Report to Westminster Government by The Wildlife Trusts.

Wallis, D.W., Hodgson, J.A. (2012) *Condatis, software to assist with the planning of habitat restoration*. [version 0.6.0, desktop]. Available from: <http://wordpress.condatis.org.uk/>

14. Appendices

Appendix A – Breakdown of Habitat Composite Region (including new habitat data created as part of the brief)

Less than N years old	Data collection method	Cumulative Area (ha)	% of LA area
5 years	Desk based	1016	12.9%
	Ground Truthed Survey	55	0.7%
	Total	1071	13.6%
10 years	Desk based	1016	12.9%
	Ground Truthed Survey	73	0.9%
	Total	1089	13.8%
15 years	Desk based	1016	12.9%
	Ground Truthed Survey	2456	31.2%
	Total	3853	49.0%
20 years	Desk based	2155	27.4%
	Ground Truthed Survey	4848	61.6%
	Total	7003	89.1%
25 years	Desk based	2155	27.4%
	Ground Truthed Survey	4848	61.6%
	Total	7003	89.1%
40 years (Upper age limit of available habitat data)	Desk based	2155	27.4%
	Ground Truthed Survey	4960	63.1%
	Total	7116	90.5%
unknown age	Desk based	0	0.0%
	Ground Truthed Survey	0	0.0%
	Total	0	0.0%

Appendix B – GIS datasets used in the generation of the NRN mapping for Cannock Chase District

Dataset	Used in	Justification	Limitations
Habitat Composite Region (including newly mapped areas as part of brief)	Strategic areas mapping. Habitat distinctiveness mapping	Provides complete coverage of the District	Wide range of ages and sources (See Appendix 2) which may limit accuracy.
OS MasterMap	Creation of new habitat polygons for Part A of the brief – Phase 1 study.	Spatial information for each field parcel, house garden etc.	No 'habitat' data within the background table data.
Land Classification data	Defining 'Pasture and Arable' and 'Urban' areas in the Habitat Connectivity Opportunity areas.	Quickly and easily define 'habitat' for large areas of land.	Very broad scale areas, covering multiple fields etc.
Functional Ecological Units	Habitat Connectivity Opportunity mapping	Only current dataset which reflects the overall areas of influence for Meres and Mosses in Staffordshire.	
Species Data (Protected Notable BAP etc.) from Staffordshire Ecological Record (SER)	Provide detail of species presence in the Habitat Connectivity Opportunity mapping.	Most complete and up-to-date database of species records in the county.	Not a consistent survey – may be some species present which are missed.
Natural Englands Priority Habitat Inventories	Strategic Areas mapping, Habitat distinctiveness mapping, Habitat Connectivity Opportunity mapping	Identification of key habitat sites within the landscape to be conserved and connected. High value sites within the Habitat distinctiveness mapping.	
Local Wildlife Sites (LWS)	Strategic Areas mapping, Habitat distinctiveness mapping, Habitat Connectivity Opportunity mapping	Identification of key habitat sites within the landscape to be conserved and connected. High value sites within the Habitat distinctiveness mapping.	
Statutory sites maps (SSSI, SAC, RAMSAR etc.)	Strategic Areas mapping, Habitat distinctiveness mapping, Habitat Connectivity Opportunity mapping	Identification of key habitat sites within the landscape to be conserved and connected. High value sites within the Habitat distinctiveness mapping.	
British Geological Survey (BGS) Soil Property Data WMS	Habitat Connectivity Opportunity mapping	Scrutiny of modelling output of condatis for production of Habitat Connectivity Opportunity Areas ensuring that HCO is	

		within the relevant soil type for that habitat based on the where habitats already exist on that soil type.	
Natural England National Character Areas (NCA)	Strategic Areas mapping, Habitat distinctiveness mapping, Habitat Connectivity Opportunity mapping	To ensure that the identified network aligns with national priorities for species, habitats and landscape.	
Staffordshire Biodiversity Action Plan (SBAP) Ecosystem Action Plan Areas (EAPS)	Habitat Connectivity Opportunity mapping	Ratification that the new Habitat Connectivity Opportunity areas are based on what has been identified as a priority in the SBAP.	

Appendix C – Evidence base confidence review (table supplied as a digital Appendix)

Staffordshire Wildlife Trust (SWT)/Staffordshire Ecological Record (SER) hold and manage the a large quantity of the county's primary ecological data which is a key factor in being able to establish a robust evidence base for any strategic environmental work. It is critical that a thorough investigation of the available datasets both in-house and those available either through Open Government Licences, a Creative Commons open licence or via a paid licence subscription to ensure that we are using the best possible datasets in the creation of the NRN.

Desirability and reliability values were scored out of 10, a list of positive and negative indicators were used to define the values for each dataset. The desirability and reliability figures were then multiplied together to give the overall 'confidence' rating which is scored out of 100, the higher the score the higher the 'confidence' of the dataset contributing to a meaningful evidence base. It must however be noted that the dataset confidence ratings are only accurate to the time that they were produced, as new datasets become available and the existing datasets are updated the confidence ratings will alter to reflect any relevant changes. The inventory therefore must be kept up to date and reviewed prior to starting any future large scale projects to ensure that the best evidence base is being used.

Appendix D – Breakdown of habitats and sites included in the habitat distinctiveness mapping bands

Distinctiveness Band	Habitats included within the band	Action (in order of preference)
Very High	<ul style="list-style-type: none"> • Irreplaceable habitats (e.g. ancient woodland) • International, national or regional value species populations. • Priority habitats as defined in Section 41 of the Natural Environment and Rural Communities (NERC) Act that are highly threatened, internationally scarce and require conservation action e.g. blanket bog 	Avoid loss, Enhance, Link, Create new habitat adjacent (expand existing habitat)
High	<ul style="list-style-type: none"> • County and District value • Habitats known to support county and District value species populations. e.g. all rivers and good quality streams. • Priority habitats as defined in Section 41 of the NERC Act requiring conservation action e.g. lowland fens 	Avoid loss, mitigate loss, last resort compensate loss. Enhance, link and create new habitat.
Medium	<ul style="list-style-type: none"> • Local Value • Habitats of Principal Importance and Staffordshire Biodiversity Action Plan (SBAP) habitats that don't meet LSW criteria, semi-natural habitats that act as corridors and stepping stones, arable land which is in a relevant stewardship agreement or organic status. • Local Value species populations. • E.g. hedges, ponds, copses and low quality woodland, rough grassland, ruderal vegetation, degraded watercourses/ditches. Habitats known to support priority species. Buildings with protected species presence that aren't high value. • Semi-natural vegetation not classed as a priority habitat e.g. hazel scrub 	Mitigate loss, compensate loss. Enhance, link and create new habitat.
Low	<ul style="list-style-type: none"> • Site Value • Intensive arable, improved and amenity grassland, manicured landscaping, isolated poor semi-natural habitat. • Semi-natural or modified vegetation not classed as a priority habitat and of lower relative value to most wildlife e.g. Temporary grass and clover ley; intensive orchard; rhododendron scrub 	Compensate large losses. Enhance, link and create new habitat.
Very Low	<ul style="list-style-type: none"> • Buildings (unless supporting protected/priority species), hard standing, roads, regularly disturbed bare ground. • Habitats and land cover of little or no value to wildlife e.g. Developed land sealed surface 	Create new habitat where connectivity exists or functional size is achievable.

Appendix E – Strategic Habitat Areas detailed methodology

The mapping works by assessing the proportion of broad habitats e.g. woodland, grassland, heathland etc. within an area to determine whether these are 'strategic', 'semi-strategic' or 'non-strategic'

Ordnance Survey 1km grid squares were classified based on the principle that if 20% or more of that square has, for instance woodland habitat within it then it is considered to function ecologically (species associated with that habitat are able to move freely within this square). Based on the above, classification of 1km squares are defined as:

- Strategic: between 5-20% of the 1km square is covered by a habitat e.g. woodland/grassland. Priority as this requires further habitat to reach the 20% threshold to be considered 'ecologically functional' for that specific habitat.
- Semi-strategic: 20% or greater specific habitat in the 1km square. Already meets the 20% threshold to be considered 'ecologically functional' but the creation of further habitat will strengthen ability for species to be able to exist and move through this square.
- Non-strategic: less than 5% of the 1km square is covered by a specific habitat making it too onerous to bring the amount of habitat to meet the 20% threshold, it is therefore not a priority area to target biodiversity compensation.

Strategic area mapping is carried out on a per habitat basis, e.g. a strategic areas map is produced for each habitat analysed, however an overall strategic areas map has been produced based on the combination of all the habitats analysed as part of the strategic mapping exercise (map 2). For this map, the methodology has been altered so that the criteria for strategic and semi-strategic areas have been swapped e.g. anything with over 20% habitat coverage is now considered strategic. By altering the methodology in this way it is possible to create a coarse overall 'connectivity map' by highlighting the areas with highest combined overall habitat availability and connectivity as opposed to those areas where it is best to create habitats.

As only higher quality habitats are assessed through this analysis (e.g. species rich grassland) and lower quality habitats are not included (table E1) (e.g. improved grassland or poor semi-improved grassland) as they do not adequately contribute to the network as they cannot support the same level of species diversity as higher quality habitats and therefore would not be able to support this diversity. This is not to say that these habitats do not contribute to the network in some way but are not presently of a high enough biodiversity value to act as a potential source site for biodiversity or to support species typical of that habitat indefinitely.

It is important to note that updating the strategic area maps over time requires up-to-date mapping data which should be sent to the Local Environmental Records Centre (LERC) when available in a suitable format to incorporate into the Nature Recovery Network Mapping.

Table E1 – Habitat types included in the assessment of strategic habitat areas (habitats without an 'X' in a relevant habitat column were not used in the assessment).

Habitat survey type	HABCODE	Habitat description	Woodland	Wetland	Grassland	Heathland
UKBAP	CF1	Coastal floodplain grazing marsh		X	X	
UKBAP	WW	Wet Woodland (Where identified)	X	X		
Phase 1	A111	Broad-leaved semi-natural woodland	X			
Phase 1	A112	Broad-leaved plantation	X			
Phase 1	A121	Coniferous semi-natural woodland	X			
Phase 1	A122	Coniferous plantation	X			
Phase 1	A131	Mixed semi-natural woodland	X			
Phase 1	A132	Mixed plantation	X			
Phase 1	A21	Dense continuous scrub	X			
Phase 1	A22	Scattered scrub	X		X	
Phase 1	A31	Broad-leaved parkland/scattered trees	X		X	
Phase 1	A32	Coniferous parkland/scattered trees	X		X	
Phase 1	A4	Recently felled woodland				
Phase 1	A5	Orchard	X		X	
Phase 1	B11	Unimproved acidic grassland			X	
Phase 1	B12	Semi-improved acidic grassland			X	
Phase 1	B21	Unimproved neutral grassland			X	
Phase 1	B22	Semi-improved neutral grassland			X	
Phase 1	B31	Unimproved calcareous grassland			X	
Phase 1	B32	Semi-improved calcareous grassland			X	
Phase 1	B4	Improved grassland				
Phase 1	B5	Marsh/marshy grassland		X	X	
Phase 1	B6	poor semi-improved grassland				
Phase 1	C11	Continuous bracken				
Phase 1	C31	Tall ruderal			X	
Phase 1	C32	Non-ruderal				
Phase 1	D11	Acid Dry dwarf shrub heath				X
Phase 1	D2	Wet dwarf shrub heath				X
Phase 1	D3	Lichen/bryophyte heath				X
Phase 1	D4	Montane heath/dwarf herb				X
Phase 1	D5	Dry heath/acidic grassland mosaic			X	X
Phase 1	D6	wet heath/acid grassland mosaic				X
Phase 1	E11	Sphagnum Bog		X		
Phase 1	E2 (any)	Flush and Spring		X	X	
Phase 1	E3 (any)	Fen		X	X	
Phase 1	F (any)	Swamp, marginal and inundation		X		
Phase 1	G (any)	Open Water		X		
Phase 1	I21	Quarry				
Phase 1	I22	Spoil				

Phase 1	I24	Refuse tip				
Phase 1	J11	Arable				
Phase 1	J112	Allotments				
Phase 1	J113	Set-aside (field margins)			X	
Phase 1	J12	Amenity grassland				
Phase 1	J13	Ephemeral/short perennial				
NVC	A (Any)	Aquatic Communities		X		
NVC	CG02	<i>Festuca ovina</i> – <i>Avenula pratensis</i> grassland			X	
NVC	CG03	<i>Bromus erectus</i> grassland			X	
NVC	CG07	<i>Festuca ovina</i> – <i>Hieracium pilosella</i> – <i>Thymus praecox/pulegioides</i> grassland			X	
NVC	H08	<i>Calluna vulgaris</i> – <i>Ulex gallii</i> heath				X
NVC	H09	<i>Calluna vulgaris</i> – <i>Deschampsia flexuosa</i> heath				X
NVC	H09/MG10	<i>Calluna vulgaris</i> – <i>Deschampsia flexuosa</i> heath / <i>Holcus lanatus</i> – <i>Juncus effusus</i> rush-pasture		X	X	X
NVC	H09/U05	<i>Calluna vulgaris</i> – <i>Deschampsia flexuosa</i> heath / <i>Nardus stricta</i> – <i>Galium saxatile</i> grassland			X	X
NVC	H09/U2	<i>Calluna vulgaris</i> – <i>Deschampsia flexuosa</i> heath / <i>Deschampsia flexuosa</i> grassland			X	X
NVC	H09a	<i>Calluna vulgaris</i> – <i>Deschampsia flexuosa</i> heath				X
NVC	H09b	<i>Calluna vulgaris</i> – <i>Deschampsia flexuosa</i> heath				X
NVC	H09c	<i>Calluna vulgaris</i> – <i>Deschampsia flexuosa</i> heath				X
NVC	H09e	<i>Calluna vulgaris</i> – <i>Deschampsia flexuosa</i> heath				X
NVC	H12	<i>Calluna vulgaris</i> – <i>Vaccinium myrtillus</i> heath				X
NVC	H12a	<i>Calluna vulgaris</i> – <i>Vaccinium myrtillus</i> heath				X
NVC	H12c	<i>Calluna vulgaris</i> – <i>Vaccinium myrtillus</i> heath				X
NVC	M22	<i>Juncus subnodulosus</i> – <i>Cirsium palustre</i> fen-meadow		X		
NVC	M23	<i>Juncus effusus/acuteiflorus</i> – <i>Galium palustre</i> rush-pasture		X		
NVC	M24	<i>Molinia caerulea</i> – <i>Cirsium dissectum</i> fen-meadow		X		
NVC	M25	<i>Molinia caerulea</i> – <i>Potentilla erecta</i> mire		X		
NVC	M26	<i>Molinia caerulea</i> – <i>Crepis paludosa</i> mire		X		
NVC	MG04	<i>Alopecurus pratensis</i> – <i>Sanguisorba officinalis</i> grassland			X	
NVC	MG05	<i>Cynosurus cristatus</i> – <i>Centaurea nigra</i> grassland			X	
NVC	MG08	<i>Cynosurus cristatus</i> – <i>Caltha palustris</i> grassland			X	
NVC	MG09	<i>Holcus lanatus</i> – <i>Deschampsia cespitosa</i> grassland			X	
NVC	MG10	<i>Holcus lanatus</i> – <i>Juncus effusus</i> rush-pasture		X	X	
NVC	S (Any)	Salt-marsh communities		X		
NVC	U01	<i>Festuca ovina</i> – <i>Agrostis capillaris</i> – <i>Rumex acetosella</i> grassland			X	
NVC	U02	<i>Deschampsia flexuosa</i> grassland			X	
NVC	U03	<i>Agrostis curtisii</i> grassland			X	
NVC	U04	<i>Festuca ovina</i> – <i>Agrostis capillaris</i> – <i>Galium saxatile</i> grassland			X	
NVC	W (any)	Woodlands and Scrub	X			

Appendix F – Condatis software technical methodology.

Condatis works by modelling a landscape of habitats as if it were an electrical circuit. A circuit board consists of a number of wires joining up resistors in combinations. When a voltage is applied to the board at one end, the current will pass through the board to the other end but the amount of current passing through each wire will vary according to the resistances it meets through each pathway. Condatis considers a landscape as analogous to a circuit board, with a source population of species being considered the voltage, the links between habitat useable by these species being the resistors, and the flow of species colonising the available habitat across those links being considered the current. Condatis is able to measure the flow of a hypothetical species across a landscape based on the availability of a distinct habitat category e.g. woodland or grassland.




Image 1 Electrical circuit on the left and comparable stylised habitat map on the right. Green represents adding a resistor or additional habitat to each to increase the number of pathways available and therefore improve the flow. Image available at: <http://wordpress.condatis.org.uk/>

Habitat source and target locations are specified: the source either representing a nominal population of species or an actual population (in this case a nominal population was used), the target representing an area for eventual colonisation. The direction of travel is defined by the placement of source and target and will depend on the purpose of study. For instance, if looking at likely species movement due to climate change, a south to north or lowland to upland direction might be required. A South-north orientation was chosen for the source and target to reflect the likely species movement change in response to climate change. Condatis looks at how the habitat in between the source and target could contribute to the species progress over multiple generations, so it is not designed to look in detail at individual patch-to-patch movements.


By using Condatis to output the relative flow of a species through the landscape for a given habitat type it is possible to more accurately define where wildlife corridors exist and where they could be improved.

Appendix G – Habitat Connectivity Opportunity areas (HCO) technical details, principles and mechanisms for delivery.


- The mapping takes into account existing local wildlife-rich habitats and existing ecological networks as well as local national and internationally designated nature conservation sites.
- The aim of the HCO areas is not to replace large areas of farmed land; we must continue to rely on working with farmers and landowners to manage existing habitats and create areas of new habitat.
- Developments whose primary objective is to conserve or enhance biodiversity particularly those which are aligned with the opportunity areas should be supported, and opportunities to incorporate biodiversity improvements in and around developments should be encouraged especially where this benefits overall biodiversity and habitat connectivity for example the creation of species rich grassland within the grassland opportunity area.
- When delivering against the mapping, care should be taken to ensure that the best possible habitat for that area is being created; it may be tempting for example where an area is both within a connectivity zone for woodland and grassland to plant large tracts of woodlands as this is easiest and most cost effective when in fact this may in some cases may result in the loss of important habitats whereas species rich grassland enhancement would be both more beneficial and provides better outcomes for habitat connectivity.
- The main aims are to ensure adequate habitats are large enough to resist harmful effects, and are well-enough connected to ensure that species are able to move around and sustain populations. Harmful effects may be localised, e.g. flooding or be much more far-reaching for example climate change. The need for more, bigger, better and joined up habitats is explained in detail in Lawton et al. (2010).
- The opportunity areas reflect and refine the work of the Staffordshire Biodiversity Action Plan Ecosystem Action Plan areas (Appendix 11) by using finer detail data to pick out more targeted conservation areas.
- The habitat connectivity opportunity areas were cross-referenced against previously mapped biodiversity opportunity zones in the District. The habitat connectivity opportunity areas are more refined than the previously mapped opportunity zones but do reflect similarities within the landscape.
- Habitat creation and restoration should take into account landscape considerations, geology and the historic environment. Particular care will be required where intensive methods are required, such as topsoil stripping / deep ploughing, or where the effect, such as woodland planting is likely to be visible from settlements or rights of way.

- Habitat creation or restoration may create opportunities too, for example screening unsightly features, creating geological exposures or helping conserve historic features.
- Regular updates of the maps is required to reflect any changes in mapped habitats as a result of physical habitat changes on the ground.


Appendix H - Previous biodiversity opportunity assessment in Cannock Chase District (2009/2010)



Appendix I – National Character Areas (NCA) in Cannock Chase District (NCA boundaries © Natural England)




Appendix J – Staffordshire Biodiversity Action Plan (SBAP) Ecosystem Action Plan Areas (EAPs) within Cannock Chase District (2019)




Appendix K – Example of how the strategic mapping will evolve over time.


1: A small gap is identified between opportunity areas for grassland (Orange shaded squares denote the opportunity area).



2: Broad scale aspirations for the creation, restoration or enhancement of species-rich grassland are identified (blue areas).



3: In time some of the aspirations are realised, leading to enhanced grassland habitat, changes monitored and mapped (orange areas).



4: The newly mapped habitat data has now influenced the opportunity area connecting two previously separate opportunity areas.

